



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 7/48, 31/00, 31/19		A1	(11) International Publication Number: WO 00/48567 (43) International Publication Date: 24 August 2000 (24.08.00)		
(21) International Application Number: PCT/US00/03902		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).			
(22) International Filing Date: 15 February 2000 (15.02.00)					
(30) Priority Data: 09/252,318 18 February 1999 (18.02.99) US					
(71) Applicant: ALLERGAN SALES, INC. [US/US]; 2525 Dupont Drive, Irvine, CA 92612 (US).					
(72) Inventor: SEFTON, John; P.O. Box 714, Trabuco Canyon, CA 92678 (US).					
(74) Agents: BARAN, Robert, J. et al.; Allergan Sales, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).					
(54) Title: METHOD AND COMPOSITION FOR TREATING ACNE					
(57) Abstract					
<p>The present invention provides a method for treating acne vulgaris by serially applying a topical composition of azelaic acid and a topical composition of benzoyl peroxide. The present invention also provides topical compositions of a peroxide of benzoyl peroxide, and azelaic acid and its derivatives, such as azelaic acid, sodium salt or methylester which are useful for treating acne vulgaris and may be used to simultaneously apply benzoyl peroxide and azelaic acid.</p>					

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHOD AND COMPOSITION FOR TREATING ACNE

FIELD OF THE INVENTION

5

This invention relates to a method and composition for treating acne vulgaris.

BACKGROUND OF THE ART

10

Acne vulgaris is an inflammatory disease of the sebaceous glands characterized by an eruption of the skin, often pustular in nature but not suppurative. Acne is a common affliction of the adolescent and affects a small but significant percentage of the adult population. Acne involvement results in unsightly lesions, particularly on the face, and in some cases results in severe scarring.

15

Various topical agents are utilized in the treatment of acne and these include sulfur, resorcinol, salicylic acid, benzoyl peroxide, retinoids and topical antibiotics. An effective anti-acne agent (or composition) must exhibit the following activities:

20

- (a) a sebostatic activity so as to inhibit hyperseborrhea;
- (b) a keratolytic and comedolytic activity so as to avoid hyperkeratosis of the follicles and to permit removal of comedos;
- 25 (c) a bacteriostatic activity so as to inhibit the activity of *Propionibacterium acnes*.

25

Nevertheless, acne vulgaris is seldom cured and only can be contained with difficulty.

30

The antibiotic clindamycin has been used, topically, to treat acne vulgaris. (See U.S. Patent 3,969,516, to Stoughton.) Various references

-2-

discuss the use of vehicle formulations to enhance the efficacy of topically-applied clindamycin. (See U.S. Patents 3,932,653; 3,989,816; 3,991,203; 4,132,781; 4,671,956; 4,746,675; 4,789,667; 4,803,228 and 4,882,359.) Clindamycin salts, clindamycin derivatives, and various dosage forms of 5 clindamycin have also been discussed as a treatment for acne vulgaris. (See U.S. Patent 3,849,396; 4,621,075 and 4,916,118.) Finally, combinations of clindamycin and other compounds active for the treatment of acne vulgaris are disclosed in U.S. Patents, 4,323,558; 4,505,896; 4,607,101; 4,906,617; 4,942,031 and 4,018,918.

10 Benzoyl peroxide has been suggested for treating acne vulgaris. (See U.S. Patent 4,387,107.) For many years, benzoyl peroxide has been proven to be a particularly powerful keratolytic and anti-seborrhic agent, as well as being endowed with antibacterial properties. Topical benzoyl peroxide compositions, including a vehicle to enhance the efficacy thereof, are 15 known (See U.S. Patent 4, 411,893). Topical compositions of benzoyl peroxide combination with antibiotics are also known. (See U.S. Patents 4,407,794; 4,692,329 and 4,387,107)

20 Peroxides, other than benzoyl peroxide, have been suggested for treatment of acne vulgaris, alone, or in combination with other compounds useful in treating acne vulgaris. (See U.S. Patents 4,607,101 and 4,906,617.) These peroxides are suggested as having certain advantages, e.g. stability over benzoyl peroxide. U.S. Patent 4,671,956 identifies the problem of benzoyl peroxide decomposing coingredients in topical formulations to thereby cause itching upon application. It is suggested that this problem 25 may be solved by including a sunscreen in the topical formulation to retard this decomposition effect of benzoyl peroxide.

Azelaic acid has been used topically and systemically to treat acne. See, for example, U.S. Patent 4,386,104 to Nazzo-Pavarro.

30 In view of the above, it is apparent that there is a great deal of interest in utilizing topical compositions for the treatment of acne vulgaris,

-3-

such compositions utilizing as an active ingredient clindamycin or benzoyl peroxide or azelaic acid, alone, or in combination with other active ingredients for the treatment of acne vulgaris.

5 Therefore, one object of the instant invention is to provide a method of treating acne vulgaris with topical compositions including benzoyl peroxide and azelaic acid.

It is another object of the invention to provide a method of treatment of acne vulgaris by topical application of compositions of benzoyl peroxide in a gel form and azelaic acid in a cream form.

10 It is another object of this invention to provide compositions for the topical treatment of acne vulgaris.

Another object of the invention is to provide topical compositions of benzoyl peroxide and azelaic acid that may be used for treating acne vulgaris.

15 Other objects and advantages of the instant invention will become apparent from a careful reading of the specification below.

SUMMARY OF THE INVENTION

20 The present invention provides a method for treating acne vulgaris by topically applying a composition of benzoyl peroxide and azelaic acid, serially, in a therapeutically-effective amount. The azelaic acid or its pharmaceutically acceptable salts or prodrugs (e.g. azelaic acid, sodium salt or lower alkyl ester), may be applied in an amount sufficient to provide from about 0.1 to about 30 percent, and preferably from about 0.5 to about 25 percent, by weight, e.g. about 20 percent azelaic acid. Benzoyl peroxide, which has keratolytic and antiseborrheic properties, may be present in an amount sufficient to provide from about 0.1 to about 30 percent, and preferably from about 2.5 to about 10 percent, by weight, benzoyl peroxide. 30 The benzoyl peroxide may more preferably be used as hydrous benzoyl

peroxide and may be suspended preferably in the form of microparticles. The above described topical composition may be in the form of a solution, gel, ointment, cream, a liquid suspension or emulsion or a stick base.

5 **DETAILED DESCRIPTION OF THE INVENTION**

The compositions of this invention are administered topically to treat acne vulgaris. That is, the compositions may be applied as a solution, gel, ointment, cream, a liquid suspension or emulsion or a stick base. Thus, it is 10 preferred that such compositions include a pharmaceutically acceptable carrier that enhances the efficacy of such topical administration. Pharmaceutically acceptable carriers include conventional emulsifiers, such as fatty alcohols, glycol ethers and esters of fatty acids; conventional 15 emollients, such as isopropyl and butyl esters of fatty acids, e.g. isopropyl myristate; humectants such as glycerin, propylene glycol, polyethylene glycol; and alcohols and acetone; oils such as mineral oil, petroleum oil, oil extracts from animal or vegetable sources; conventional stabilizers including antioxidants and preservatives. The compositions may also 20 include agents, such as urea, to improve the hydration of the skin. In addition to the foregoing conventional formulations, the topical 25 compositions may include penetration-enhancing agents such as 1-pyrrolidone and N-lower alkyl-2-pyrrolidones, such as N-methyl-2-pyrrolidone; and 1-substituted azacycloalkan-2-ones such as, for example, 1-n-dodecylazacycloheptan-2-one and other compounds disclosed in U.S. Pat. No. 3,989,816. Longer chain sulfoxides, e.g., n-octyl methyl sulfoxide and hexamethylene-lauramide and the other penetration-enhancing agents disclosed in U.S. Patent No. 4,743,588, may also be included in the formulations utilized in the method of this invention. The amount of these penetration-enhancing agents which may be used in the present invention

ranges from about 0.1 to 25 percent and preferably about 1 to 15 percent by weight of the composition.

5 The amount of the compositions to be administered will obviously be an effective amount for the desired result expected therefrom. This, of course, will be ascertained by the ordinary skill of the practitioner. In accordance with the usual prudent formulating practices, a dosage near the lower end of the useful range of the particular agent may be employed initially and the dosage increased as indicated from the observed response, as in the routine procedure of the physician.

10 In carrying out the novel method employing the topical route, the active ingredient(s) formulated, for example, as a gel or lotion or suspension, is applied to the affected area of the skin at a rate varying from 0.2 mg per square cm of skin surface per day up to 10 mg per square cm of skin surface per day until the appearance of the affected skin has returned 15 to normal. The gel or lotion or suspension is generally applied for several days.

The topical compositions of this invention may be applied to the face of a patient with acne 1 to 4 times daily with the result that open and closed comedones are markedly reduced within two to four weeks.

20 The topical composition, including azaleic acid, is preferably a cream formulation. Typically, said cream formulation may comprise:

FORMULATION A

<u>INGREDIENTS</u>	<u>WEIGHT PERCENT</u>
AZELAIC ACID	20.00
BENZOIC ACID DAB	0.20
PROPYLENE GLYCOL USP	12.50
CUTINA CBS (1)	7.00
PEG-5 GLYCERYL STEARATE (2)	5.00

-6-

CETEARYL OCTANOATE (3)	3.00
GLYCERIN (85%) DAB	1.5
PURIFIED WATER DAB (4)	QS

5 1. Glyceryl Stearate (and) Cetearyl Alcohol (and) Cetyl Palmitate (and)
 Cocoglycerides (HENKEL)
2. Arlatone 983S (ICI)
3. PCL Liquid (DRAGOCO)
4. 50.80 percent (w/w)

10

The topical composition, including benzoyl peroxide, is preferably a gel formulation. Typically, said gel formulation may comprise:

FORMULATION B

15

4 or 8 percent, by weight, benzyl peroxide in a gel vehicle containing purified water, cetyl alcohol dimethyl isosorbide, fragrance, simethicone, stearyl alcohol and ceteareth-20.

20 An example of such a product is Brevoxyl® benzyl peroxide available from Stiefel Laboratories, Inc., Coral Gables, FL.

The invention is further illustrated by the following formulations and examples which are illustrative of a specific mode of practicing the invention and is not intended as limiting the scope of the claims.

25

EXAMPLE 1

A 20 year old male applies 0.35 gms of the Formulation A and 0.35 gms of Formulation B to his face 4 times daily, each. After 10 days, the number of comedones begin to diminish. By the end of four weeks, the 30 number of comedones declines significantly.

EXAMPLE 2

29 patients were treated, serially, with Formulation A and Formulation B as compared to a control group of 29 patients which was treated with Benzamycin® acne medication, comprising benzyl peroxide 5% and erythromycin 3%, by weight, available from Dermik Laboratories, Inc., Collegeville, PA 19426. 25 patients from each group completed the study. The demographic data for these two groups are reported in Table 1, below.

Table 1
Demographic Data

		Azelex/BP			Benzamycin		
		Male	Female	Total	Male	Female	Total
Age: n	11 (37.9%)	18 (62.1%)	29	5 (17.2%)	24 (82.8%)	29	29
	20.6 ^a	28.9 ^a	25.8	20.7 ^b	30.7 ^b	28.9	28.9
	8.37	6.80	8.37	4.72	5.99	6.89	6.89
	13.3	18.8	13.3	15.0	19.4	15.0	15.0
	39.6	39.5	39.6	25.5	44.0	44.0	44.0
Race: n	5 (45.4%)	10 (55.6%)	15 (51.7%)	0	15 (62.5%)	15 (51.7%)	15 (51.7%)
	6 (54.6%)	7 (38.9%)	13 (44.8%)	5 (100%)	9 (37.5%)	14 (48.3%)	14 (48.3%)
	0	0	0	0	0	0	0
	0	1 (5.6%)	1 (3.4%)	0	0	0	0
	0	0	0	0	0	0	0
	0	0	0	0	0	0	0
Acne Hx: n	< 1 year	0	0	0	0	0	0
	1-2 years	6 (54.6%)	1 (5.6%)	7 (24.1%)	2 (40.0%)	0	2 (6.9%)
	3-5 years	1 (9.1%)	3 (16.7%)	4 (13.8%)	2 (40.0%)	2 (8.3%)	4 (13.8%)
	6-10 years	1 (9.1%)	5 (27.8%)	6 (20.7%)	1 (20.0%)	9 (37.5%)	10 (34.5%)
	> 10 years	3 (27.3%)	9 (50.0%)	12 (41.4%)	(%)	13 (54.2%)	13 (44.8%)

^a = p < 0.05, within Azelex/BP group, for age between, male - female

^b = p < 0.05, within Benzamycin group, for age between, male - female

^c = p < 0.05, within Benzamycin group, for race between, male - female

^d = p < 0.05, within Benzamycin group, for acne history between, male - female

-9-

The patients were first evaluated for the severity of their acne conditions. This evaluation is reported in Table 2, below.

-12-

TABLE 3

Overall Disease Severity-Improvement From Baseline

5

10

		<u>Week 4</u>	<u>Week 8</u>	<u>Week 12</u>
	Azelex/BPO	1.3	1.5	1.8
	Benzamycin	0.9	0.7	1.0

-13-

Table 4
Lesion Count - Pustules

Visit Number	Treatment Group	Change from Baseline (Week 0)							
		n	mean	SD	p-value	n	mean	SD	p-value
1	Azealex/BP	29	5.4	6.59	0.566				
(Week 0)	Benzamycin	29	4.1	5.70					
2	Azealex/BP	24	1.9	3.15	0.092	24	3.6	5.95	0.329
(Week 4)	Benzamycin	24	2.6	2.86		24	2.0	5.15	
3	Azealex/BP	22	1.4	3.24	0.004	22	4.6	7.44	0.313
(Week 8)	Benzamycin	19	2.0	1.81		19	2.4	6.25	
4	Azealex/BP	25	1.6	3.43	0.553	25	3.8	7.70	0.141
(Week 12)	Benzamycin	25	2.7	7.47		25	1.2	3.72	

As reported in Tables 3 and 4, above, the serial treatment with azaleic acid and benzoyl peroxide is significantly more effective for treating acne than the Benzamycin® control.

Similar evaluations for open comedones, closed comedones and inflammatory lesions show that the method of this invention is more effective than the control.

As to the side effects of treating patients for acne by the method of the invention and the control, the following results were obtained.

10

Scaling.	Azaleic acid (AA)/Benzylperoxide (BPO) is slightly better than the control.
Erythema.	AA/BPO is better than the control.
Dryness.	AA/BPO is better than the control.
Oiliness.	AA/BPO is equal to or slightly worse than the control.
Burning.	AA/BPO is better than the control.
Itching.	AA/BPO is better than the control.

15

While particular embodiments of the invention have been described, it will be understood, of course, that the invention is not limited thereto since many obvious modifications can be made; and it is intended to include within this invention any such modifications as will fall within the scope of the appended claims. For example, it will be appreciated by those skilled in the art that various pharmaceutically acceptable derivatives, salts and prodrugs of azelaic acid, e.g. azelaic acid, sodium or potassium salt, or lower alkyl ester, i.e. C₁ to C₆ alkyl ester, e.g. methyl azelate, may be used in place of azaleic acid. Also, various forms of peroxides may be used in place of hydrous benzoyl peroxide; i.e., diaryl peroxide, alkyl aryl peroxide, cycloalkyl aryl peroxide, may be substituted for hydrous benzoyl peroxide. For example, lauroyl benzoyl peroxide, cyclohexyl carbanoyl benzoyl peroxide may be used in place of benzoyl peroxide.

20

Also, the azaleic acid and benzyl peroxide may be combined in a single topical composition, e.g., a cream or gel, for ease of application. Typically, said single topical composition will comprise the amounts of

-15-

azaleic acid and benzoyl peroxide sufficient to provide the amounts described above for serial application in a single topical application.

CLAIMS:

1. A method of treating acne vulgaris in human patients comprising topically administering to said patients separate compositions comprising a therapeutically effective amount of benzoyl peroxide and azelaic acid, in a serial manner.
5
2. The method of claim 1 wherein said benzoyl peroxide and azelaic acid are dispersed in pharmaceutically acceptable carriers.
10
3. The method of claim 2 wherein said composition is a cream or a gel.
15
4. The method of claim 3 wherein one of said compositions comprises from about 0.1 to 30 percent, by weight, azelaic acid.
15
5. The method of claim 4, wherein one of said compositions comprises from about 0.1 to 30 percent, by weight, benzoyl peroxide.
20
6. The method of claim 5 wherein said azaleic acid composition comprises a cream formulation.
25
7. The method of claim 6 wherein said benzyl peroxide composition comprises a gel composition.
25
8. The method of claim 7 wherein said composition comprises from about 0.5 to 25 percent, by weight azelaic acid.
30
9. The method of claim 8 wherein said composition comprises about 20 percent, by weight, azelaic acid.
30

-17-

10. The method of claim 9, wherein said composition comprises from about 2.5 to 10 percent, by weight, benzyl peroxide.

5 11. A topical composition for treating acne vulgaris in human patients comprising a therapeutically effective amount of benzoyl peroxide and azelaic acid.

10 12. The composition of claim 11 wherein said benzoyl peroxide and azelaic acid are dispersed in a pharmaceutically acceptable carrier.

13. The composition of claim 12 wherein said composition is a cream or gel.

15 14. The composition of claim 13 wherein said composition comprises from about 0.1 to 30 percent, by weight, azelaic acid.

16. The composition of claim 14 wherein said composition comprises from about 2.5 to 10 percent, by weight, benzoyl peroxide.

20 16. The composition of claim 15 wherein said composition comprises about 0.5 to 25 percent, by weight, azelaic acid.

25 17. The composition of claim 16 wherein said composition comprises about 20 percent, by weight, azelaic acid.

18. A method of treating acne vulgaris in human patients which comprises topically administering to said patients a composition according to claim 11.

30

-18-

19. A method of treating acne vulgaris in human patients which comprises topically administering to said patients a composition according to claim 14.

5 20. A method of treating acne vulgaris in human patients which comprises topically administering to said patients a composition according to claim 17.

INTERNATIONAL SEARCH REPORT

Inte
nal Application No
PCT/US 00/03902

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K7/48 A61K31/00 A61K31/19

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, WPI Data, PAJ, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category ^o	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GB 2 150 436 A (L'OREAL) 3 July 1985 (1985-07-03) examples 1-8 & US 4 671 956 A 9 June 1987 (1987-06-09) cited in the application -----	1-20
Y	US 4 386 104 A (NAZZARO-PORRO) 31 May 1983 (1983-05-31) cited in the application claims 1,4,5; example 3 -----	1-20
X	DE 43 07 976 A (BEIERSDORF AG) 22 September 1994 (1994-09-22) page 2, line 66 -page 3, line 2 page 3, line 8 examples 1,5 -----	11-17

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

^o Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

10 July 2000

Date of mailing of the international search report

14/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alvarez Alvarez, C

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 00/03902

Patent document cited in search report	Publication date	Patent family member(s)			Publication date
GB 2150436	A 03-07-1985	LU	85111 A		12-09-1985
		AT	390372 B		25-04-1990
		AT	380384 A		15-10-1989
		AU	567597 B		26-11-1987
		AU	3606284 A		06-06-1985
		BE	901171 A		30-05-1985
		CA	1220717 A		21-04-1987
		CH	660687 A		15-06-1987
		DE	3443647 A		13-06-1985
		FR	2555898 A		07-06-1985
		IT	1178653 B		16-09-1987
		JP	4045491 B		27-07-1992
		JP	60142920 A		29-07-1985
		NL	8403659 A		01-07-1985
		US	4671956 A		09-06-1987
US 4386104	A 31-05-1983	IT	1075741 B		22-04-1985
		IT	1114950 B		03-02-1986
		AU	524602 B		23-09-1982
		AU	3502778 A		18-10-1979
		BE	866151 A		19-10-1978
		CA	1137873 A		21-12-1982
		CH	638099 A		15-09-1983
		DE	2817133 A		02-11-1978
		GB	1603563 A		25-11-1981
		JP	1455083 C		25-08-1988
		JP	53130433 A		14-11-1978
		JP	63001287 B		12-01-1988
		ZA	7802125 A		28-03-1979
		US	4818768 A		04-04-1989
		US	4292326 A		29-09-1981
DE 4307976	A 22-09-1994	WO	9421241 A		29-09-1994
		EP	0689431 A		03-01-1996
		JP	8507772 T		20-08-1996